

Common Repatha® Documentation Requirements for Patients With Primary Hyperlipidemia and Established CVD^{1,2}

Information Category	Clinical Details	Documentation Examples
Primary Diagnosis	<input type="checkbox"/> Primary hyperlipidemia	ICD-10 code*: <ul style="list-style-type: none"> • E78.00 Pure hypercholesterolemia, unspecified • E78.2 Mixed hyperlipidemia • E78.4 Other hyperlipidemia • E78.5 Hyperlipidemia, unspecified
Secondary Diagnosis	<input type="checkbox"/> Established CVD, including but not limited to: <ul style="list-style-type: none"> • Acute coronary syndrome • History of myocardial infarction • Stable or unstable angina • Coronary or other arterial revascularization • Stroke • Peripheral arterial disease (PAD) 	Select the appropriate secondary ICD-10 code* [†] (examples of ICD-10 codes can be found below or on the RepathaReady® or Specialty Pharmacy Intake Form)
Diagnostic Findings	<input type="checkbox"/> Baseline and current lipid levels <input type="checkbox"/> Diagnostic tests	<ul style="list-style-type: none"> • Lipid level values and lab dates (at baseline and most recent) <ul style="list-style-type: none"> - Low-density lipoprotein cholesterol (LDL-C) - Triglyceride - Total cholesterol • Diagnostic test results and dates (angiogram, stress test, echocardiogram, nuclear imaging, etc) • CVD risk score (Framingham score, Pooled Cohort Equations calculation, etc)
Experience With Repatha®	<input type="checkbox"/> Repatha®	<ul style="list-style-type: none"> • Percentage reduction in LDL-C level (goal, achieved, etc) • Duration/dates of therapy • LDL-C level pre- and post-treatment
History of Other Lipid-lowering Treatment <i>(may include multiple therapies)</i>	<input type="checkbox"/> High-intensity statin: <ul style="list-style-type: none"> • Atorvastatin 40 mg or 80 mg • Rosuvastatin 20 mg or 40 mg <input type="checkbox"/> Other therapies (eg, ezetimibe)	<ul style="list-style-type: none"> • Duration/dates of therapy • LDL-C level pre- and post-treatment

OR Intolerance/Contraindication to Other Lipid-lowering Treatments

Intolerance or Contraindication to Other Lipid-lowering Treatment	<input type="checkbox"/> Intolerance, including but not limited to: <ul style="list-style-type: none"> • Rhabdomyolysis • Muscle pain or weakness • Elevated creatine kinase (CK) • Elevated liver function tests <input type="checkbox"/> Contraindication <ul style="list-style-type: none"> • Contraindication to treatment 	<ul style="list-style-type: none"> • Intolerance <ul style="list-style-type: none"> - Maximum tolerated statin dose - Documented lab results - Duration of symptoms - Response to statin re-challenge • Contraindication <ul style="list-style-type: none"> - Documented contraindication
Lifestyle Modification	<input type="checkbox"/> Diet <input type="checkbox"/> Exercise	<ul style="list-style-type: none"> • Low-fat diet (documented dietitian visits, group support, etc) • Exercise (gym attendance, documented physical activity, etc) • Enrollment in weight management programs

CVD = cardiovascular disease; LDL-C = low-density lipoprotein cholesterol; PA = prior authorization.

* The sample diagnosis codes are informational and not intended to be directive or a guarantee of reimbursement, and include potential codes that would include FDA-approved indications for Repatha®. Other codes may be more appropriate given internal system guidelines, payer requirements, practice patterns, and the services rendered.

[†] Examples of secondary ICD-10 codes for established CVD*:

I20.0__ Unstable angina	I65.___ Occlusion and stenosis of cerebral arteries, extracranial
I20.9__ Angina pectoris, unspecified	I66.___ Occlusion and stenosis of cerebral arteries, intracranial
I21.___ Acute myocardial infarction	I67.___ Other cerebrovascular disease
I22.___ Subsequent myocardial infarction	I70.___ Atherosclerosis
I25.___ Chronic ischemic heart disease	I73.9__ Peripheral vascular disease, unspecified
I63.___ Cerebral infarction	G45.9__ Transient cerebral ischemic attack, unspecified
	G46.___ Vascular syndromes

Reference: 1. Data on file, Amgen; 2016. 2. Repatha® (evolocumab) Prescribing Information, Amgen.

Please see Indications and Important Safety Information on the next page.



Indications and Important Safety Information

Indications

Prevention of Cardiovascular Events: In adults with established cardiovascular disease, Repatha® is indicated to reduce the risk of myocardial infarction, stroke, and coronary revascularization.

Primary Hyperlipidemia (including Heterozygous Familial Hypercholesterolemia): Repatha® is indicated as an adjunct to diet, alone or in combination with other lipid-lowering therapies (e.g., statins, ezetimibe), for the treatment of adults with primary hyperlipidemia to reduce low-density lipoprotein cholesterol (LDL-C).

Homozygous Familial Hypercholesterolemia: Repatha® is indicated as an adjunct to diet and other LDL-lowering therapies (e.g., statins, ezetimibe, LDL apheresis) for the treatment of patients with homozygous familial hypercholesterolemia (HoFH) who require additional lowering of LDL-C.

The safety and effectiveness of Repatha® have not been established in pediatric patients with HoFH who are younger than 13 years old or in pediatric patients with primary hyperlipidemia or HeFH.

Important Safety Information

Contraindication: Repatha® is contraindicated in patients with a history of a serious hypersensitivity reaction to Repatha®. Serious hypersensitivity reactions including angioedema have occurred in patients treated with Repatha®.

Allergic Reactions: Hypersensitivity reactions (e.g. angioedema, rash, urticaria) have been reported in patients treated with Repatha®, including some that led to discontinuation of therapy. If signs or symptoms of serious allergic reactions occur, discontinue treatment with Repatha®, treat according to the standard of care, and monitor until signs and symptoms resolve.

Adverse Reactions in Primary Hyperlipidemia (including HeFH): The most common adverse reactions (>5% of patients treated with Repatha® and occurring more frequently than placebo) were: nasopharyngitis, upper respiratory tract infection, influenza, back pain, and injection site reactions.

From a pool of the 52-week trial and seven 12-week trials: Local injection site reactions occurred in 3.2% and 3.0% of Repatha®-treated and placebo-treated patients, respectively. The most common injection site reactions were erythema, pain, and bruising.

Allergic reactions occurred in 5.1% and 4.7% of Repatha®-treated and placebo-treated patients, respectively. The most common allergic reactions were rash (1.0% versus 0.5% for Repatha® and placebo, respectively), eczema (0.4% versus 0.2%), erythema (0.4% versus 0.2%), and urticaria (0.4% versus 0.1%).

Adverse Reactions in the Cardiovascular Outcomes Trial: The most common adverse reactions (>5% of patients treated with Repatha® and occurring more frequently than placebo) were: diabetes mellitus (8.8% Repatha®, 8.2% placebo), nasopharyngitis (7.8% Repatha®, 7.4% placebo), and upper respiratory tract infection (5.1% Repatha®, 4.8% placebo).

Among the 16,676 patients without diabetes mellitus at baseline, the incidence of new-onset diabetes mellitus during the trial was 8.1% in patients assigned to Repatha® compared with 7.7% in those assigned to placebo.

Adverse Reactions in Homozygous Familial Hypercholesterolemia (HoFH): The adverse reactions that occurred in at least two patients treated with Repatha® and more frequently than placebo were: upper respiratory tract infection, influenza, gastroenteritis, and nasopharyngitis.

Immunogenicity: Repatha® is a human monoclonal antibody. As with all therapeutic proteins, there is potential for immunogenicity with Repatha®.

Please [click here](#) to see full Prescribing Information.



Amgen
One Amgen Center Drive
Thousand Oaks, CA 91320-1799
www.amgen.com

© 2018 Amgen Inc. All rights reserved. USA-145-80785 12-18

